



North Shore Gastroenterology Associates, P.C.
233 East Shore Road, Suite 101
Great Neck, New York 11023
Tel: 516-487-2444
Fax: 516-487-2446
www.northshoregastro.com

**COLONOSCOPY
CLENPIQ PREP**

PATIENT NAME: _____ DATE OF PROCEDURE: _____

DOCTOR: _____ LOCATION: _____

ARRIVAL TIME: _____ PROCEDURE TIME: _____

HOLD THE FOLLOWING MEDICATIONS: _____

IF YOU ARE UNABLE TO KEEP YOUR APPOINTMENT, WE ASK THAT YOU GIVE US 72 HOURS NOTICE. If you do not call, and fail to show up for your scheduled procedure you will be charged a "No Show Fee" of \$100.00.

You MUST make arrangements to have someone accompany you to your procedure. THIS PERSON MUST COME UPSTAIRS TO SIGN YOU OUT. Due to the effects of anesthesia, YOU ARE NOT ALLOWED TO DRIVE FOR THE REST OF THE DAY. For your safety we CANNOT allow you to take a taxi.

You may NOT go to work on the day of your procedure. You MUST stay home and rest for the remainder of the day.

****IMPORTANT: IF YOU ARE A SMOKER, WE STRONGLY ADVISE THAT YOU DO NOT SMOKE THE DAY OF YOUR PROCEDURE. SMOKING PRIOR TO YOUR PROCEDURE COULD CAUSE COMPLICATIONS WITH ANESTHESIA****

PICK UP FROM THE PHARMACY:

1. Clenpiq; this is available by prescription only. Please contact our office immediately if you did not receive your prescription.

AT LEAST ONE WEEK PRIOR TO YOUR PROCEDURE:

1. **If you take any blood thinners, such as Coumadin, Plavix or Aspirin on a daily basis, you must notify your Doctor IMMEDIATELY!** These medications **MAY** need to be discontinued prior to your procedure.
2. **If you take Insulin, you must discuss with your Doctor whether or not you need to hold off taking or reduce your morning dose the day of the procedure.**
3. **STOP** taking any vitamins containing iron and/or iron supplements **ONE WEEK** prior to your procedure.

THE DAY BEFORE YOUR PROCEDURE:

Do not apply any lotion to your chest, arms or legs!

1. **You must be on a CLEAR LIQUID DIET for the ENTIRE day.**

You May Have:	You May NOT Have:
Water	Solid Foods
Seltzer	Alcohol
Apple Juice	Lemonade
Sprite	Milk or Milk Products
Ginger Ale	
Plain Jello (not red or purple)	
Bouillon or Clear Broth	
Gatorade (not red or purple)	
White Grape Juice	
Italian Ice(not red or purple)	
Black Coffee or Tea (no milk or creamer)	
Gum	

2. **At 5PM**, drink ONE (1) bottle of CLENPIQ. Drink CLENPIQ directly from the bottle. No mixing or diluting is required. **(See Illustration on next page.)** Drink five (5) 8-ounce glasses of clear liquid over the next 5 hours.
3. **You are encouraged to continue to drink as much clear liquid as possible until bedtime.**

THE DAY OF THE PROCEDURE:

1. At _____ **(5 hours before your procedure)** drink the remaining bottle of CLENPIQ. Drink CLENPIQ directly from the bottle. No mixing or diluting is required. **(See Illustration on next page.)** Drink three (3) 8-ounce glasses of clear liquid. **(See Illustration on next page.)** This must be completed by _____ **(4 hours prior to your Colonoscopy).**

IMPORTANT: YOU CANNOT HAVE ANYTHING ELSE TO EAT OR DRINK, NOT EVEN WATER, UNTIL AFTER YOUR PROCEDURE. IF YOU DO, YOUR PROCEDURE WILL BE DELAYED OR CANCELLED!

2. **YOU SHOULD TAKE YOUR BLOOD PRESSURE AND/OR HEART MEDICATIONS EARLY IN THE MORNING WITH A SMALL SIP OF WATER. If you must take other medications on the day of the procedure, please call our office.**
3. **HAVE NOTHING ELSE TO EAT OR DRINK, NOT EVEN WATER,** until after your procedure.
4. **FEMALE PATIENTS:** Please be aware that on the day of your procedure, you may need to provide a urine sample.

***** REMEMBER *****

DO NOT OMIT ANY PART OF THE PREPARATION. A POOR PREPARATION MAY CAUSE A POLYP OR CANCER TO GO UNDETECTED AND THE PROCEDURE MAY NEED TO BE REPEATED AFTER REPREPPING.

Stay hydrated: A colonoscopy prep causes the body to lose a significant amount of fluid and can result in sickness due to dehydration. It's important that you prepare your body by drinking extra clear liquids before the prep, stay hydrated by drinking all required clear liquids during the prep, and replenish your system by drinking clear liquids after returning home from your colonoscopy.

Drink extra liquids before, during, and after your bowel prep (until 4 hours before your procedure), as well as after your colonoscopy!!

Stay positive: We realize that the colonoscopy prep may not be the most glamorous process. But remember, when you get a colonoscopy, you are taking a proactive step toward prevention or early detection of colorectal cancer.